

HZPP-9802
Jan 25, 1998

The Nonlinear Spatial Damping rate in QGP *

Chen Jisheng[†] Li Jiarong

Institute of particle physics, Central China Normal University

Wuhan, 430079, Hubei,P.R. China

Abstract

The derivative expansion method has been used to solve the semiclassical kinetic equations of quark-gluon plasma. The nonlinear spatial damping rate, the imaginary part of the wave vector, for the longitudinal secondary color waves in the long wavelength limit has been calculated numerically.

Typeset using REVTeX

*Work supported by the National Natural Science Fund of China.

[†]Permanent address:University of Hydraulic & Electric Engineering/Yichang

Keywords:QGP, nonlinear spatial damping rate, derivative expansion

The Landau damping as collisionless damping is an important collective effect in quark-gluon plasma(QGP); it describes how the color field is affected by media during its traveling in QGP. In the frame of kinetic theory, it's shown that there is no Landau damping in linear or Abelian dominance approximation ^[1-3] but there is nonlinear Landau damping ^[4]. However, all the previous works give only the temporal damping rate, i.e., the instabilities with complex frequency and real wave vector values. As in the electromagnetic plasma ^[5], if the imaginary part of wave vector for the secondary waves resulted from the nonlinear interaction in plasmas may not be zero, there may be spatial damping also in QGP. The collisionless damping in QGP can be completely recognized only when the temporal and spatial damping are known. Up to now, neither the QGP kinetic theory nor the finite temperature QCD has given the spatial damping rate directly. In this letter, we will work from the QGP kinetic equations and derive the nonlinear spatial damping rate in QGP finally.

The collisionless kinetic equations for QGP are ^[6]

$$\begin{aligned} p^\mu D_\mu f(x, p) + \frac{1}{2} p^\mu \frac{\partial}{\partial p_\nu} \{F_{\mu\nu}(x), f(x, p)\} &= 0; \\ p^\mu D_\mu \bar{f}(x, p) - \frac{1}{2} p^\mu \frac{\partial}{\partial p_\nu} \{F_{\mu\nu}(x), \bar{f}(x, p)\} &= 0; \\ p^\mu \tilde{D}_\mu G(x, p) + \frac{1}{2} p^\mu \frac{\partial}{\partial p_\nu} \{\mathcal{F}_{\mu\nu}(x), G(x, p)\} &= 0 \end{aligned} \quad (1)$$

where f , $\bar{f}$, G are the distribution functions of quarks, antiquarks, gluons in QGP, respectively. $F_{\mu\nu}$, $\mathcal{F}_{\mu\nu}$ represent the mean field stress tensors in fundamental and adjoint representation, i.e., $F_{\mu\nu} = F_{\mu\nu}^a I_a$, $\mathcal{F}_{\mu\nu} = F_{\mu\nu}^a F_a$. I_a and F_a are the corresponding generators.

The mean color field equation coupled with the kinetic equations is

$$D_\mu F^{\mu\nu}(x) = j^\nu(x) \quad (2)$$

j^ν is the color current

$$j^\nu(x) = -\frac{g}{2} \int \frac{d^3p}{(2\pi)^3 E_p} p^\nu \left[(f(x, p) - \bar{f}(x, p)) + 2i I_a f_{abc} G_{bc}(x, p) \right] \quad (3)$$

where f_{abc} is the structure constant of $SU(N_c)$.

To calculate the imaginary part of the wave vector for the color field conveniently, we will solve the QGP kinetic equations iteratively in momentum space. Noting that the energy density of the color field in QGP is much smaller than the hot energy density in high temperature and density condition, the mean field strength may be selected as the separation of these scales. The derivative expansion method is an effective tool to solve the nonlinear equation [7,8]. In this method, it's essential that not only the function but also the relevant derivative in equation will be expanded iteratively. To discuss the spatial damping, we will only expand the space derivative in momentum space; the wave vector and the relevant functions can be expanded as

$$\begin{aligned}\mathbf{k} &= \sum_{j=0}^N \alpha^j \mathbf{k}^{(j)}; \\ A_i &= \sum_{j=1}^N \alpha^j A_i^{(j)}(\mathbf{k}^{(0)}, \mathbf{k}^{(1)}, \dots, \mathbf{k}^{(N)}); \\ f &= \sum_{j=0}^N \alpha^j f^{(j)}(\mathbf{k}^{(0)}, \mathbf{k}^{(1)}, \dots, \mathbf{k}^{(N)})\end{aligned}\tag{4}$$

where α is a dimensionless parameter introduced to denote the order of a small quantity. $\bar{f}$ and G can be expanded similarly. For convenience we work in the temporal gauge, $A_a^0 = 0$. The relation between the color electrical field $\mathbf{E}$ and the color vector potential $\mathbf{A}$ can be expressed as: $E_a^i = -\partial A_a^i(x)/\partial t$.

The color current at any order is expanded as

$$j^{h(n)} = -\frac{g}{2} \int \frac{d^3 p}{E_p (2\pi)^3} p^h \left\{ \left(f^{(n)} - \bar{f}^{(n)} \right) + 2i I_a f_{abc} G_{bc}^{(n)} \right\}\tag{5}$$

From the kinetic equations we can see that the distribution functions will fluctuate when the mean field is applied to a plasma system as an external field, and the fluctuating functions reversely influence the mean field. This self-consistent relation between the mean field and the distribution functions of the plasma particles are well expressed by the expansions Eqs.(4) and Eq.(5). The leading terms describe the linear approximation. The important dispersion relation obtained in the linear approximation will be changed by the nonlinear interactions

of the eigenwaves, i.e., wave vector corrections $\mathbf{k}^{(1)}$, $\mathbf{k}^{(2)}$, , $\dots$, $\mathbf{k}^{(N)}$ will be given to the eigenwave vector $\mathbf{k}^{(0)}$ correspondingly. As in electromagnetic plasma, if the imaginary parts of the wave vector correction for eigenmodes are not vanishing, there will be spatial damping.

To simplify the calculation, we suppose that there's only longitudinal vector potential:

$$A^i(k) = k^{(0)i} A(k) / K^{(0)} \quad (6)$$

where $K^{(0)} = |\mathbf{k}^{(0)}|$.

Inserting the expansions Eqs.(4) and Eqs.(5) into the equations (1-2) in momentum space, then equating the coefficients of equal power in α in two sides of these equations, one can obtain a hierarchy of equations.

The first order mean field equation is

$$-\omega^2 A^{(1)h}(k) = j^{(1)h}(k) \quad (7)$$

The first order transport equation for the distribution function of quarks is

$$\begin{aligned} p \cdot k^{(0)} f^{(1)}(k, p) + \frac{g}{2} \sum_{k_1+k_2=k} (p \cdot k_1^{(0)}) \{A_i^{(1)}(k_1), \partial_p^i f^{(0)}(k_2, p)\} \\ + \frac{g}{2} \sum_{k_1+k_2=k} p_i \{A_i^{(1)}(k_1), k_{1\nu}^{(0)} \cdot \partial_p^\nu f^{(0)}(k_2, p)\} = 0 \end{aligned} \quad (8)$$

where we have defined $k^{(0)} = (\omega, \mathbf{k}^{(0)})$ and

$$\sum_{k_1+k_2=k} = \int \frac{d^4 k_1 d^4 k_2}{(2\pi)^4} \delta^4(k - k_1 - k_2). \quad (9)$$

The first order kinetic equations for antiquarks and gluons are similar to Eq.(8) except the opposite signs of the terms related to $\{\dots, \dots\}$ for antiquarks, f and A are replaced by G and $\mathcal{A}$ for gluons, respectively.

Assuming background configuration is local neutral, the zero order colorless distribution functions can be chosen as the Fermi-Dirac and Bose-Einstein equilibrium distribution functions, respectively:

$$f^{(0)}(k, p) = \bar{f}^{(0)}(k, p) = (e^{\beta p \cdot U} + 1)^{-1}; \quad G^{(0)}(k, p) = (e^{\beta p \cdot U} - 1)^{-1} \quad (10)$$

where $\beta = 1/T$ is the inverse of the temperature and U is the local flow velocity(normalized to $U_\mu U^\mu = 1$). Using Eq.(8) and Eq.(6), we can express the first order distribution functions in terms of the corresponding zero order distribution functions and the first order field potential

$$\begin{aligned} f^{(1)}(k, p) &= -g\omega \frac{p^0 k^{(0)i}}{p \cdot k^{(0)} + ip^0 0^+} \frac{df^{(0)}(p)}{dp^i} \frac{A^{(1)}(k)}{K^{(0)}}, \\ \bar{f}^{(1)}(k, p) &= g\omega \frac{p^0 k^{(0)i}}{p \cdot k^{(0)} + ip^0 0^+} \frac{d\bar{f}^{(0)}(p)}{dp^i} \frac{A^{(1)}(k)}{K^{(0)}}, \\ G^{(1)}(k, p) &= -g\omega \frac{p^0 k^{(0)i}}{p \cdot k^{(0)} + ip^0 0^+} \frac{dG^{(0)}(p)}{dp^i} \frac{\mathcal{A}^{(1)}(k)}{K^{(0)}} \end{aligned} \quad (11)$$

By inserting Eq.(11) into the mean field equation Eq.(7) we can obtain the dispersion relation satisfied by the frequency and wave vector of the eigenwave in the first order approximation.

$$\epsilon(\omega, \mathbf{k}^{(0)}) = 1 + \frac{3\omega_p^2}{K^{(0)2}} \left[1 - \frac{\omega}{2K^{(0)}} \left(\ln \left| \frac{K^{(0)} + \omega}{K^{(0)} - \omega} \right| - i\pi\theta(K^{(0)} - \omega) \right) \right] = 0 \quad (12)$$

and the solution is

$$A_{\mathbf{k}^{(0)}}^{(1)\sigma} = -i\frac{\pi}{\omega} E_{\mathbf{k}^{(0)}}^\sigma \left[e^{-i\phi_{\mathbf{k}^{(0)}}^\sigma} \delta(\omega - \omega_{\mathbf{k}^{(0)}}^\sigma) + e^{i\phi_{\mathbf{k}^{(0)}}^\sigma} \delta(\omega + \omega_{\mathbf{k}^{(0)}}^\sigma) \right] \quad (13)$$

where $E_{\mathbf{k}^{(0)}}^\sigma$ and $\phi_{\mathbf{k}^{(0)}}^\sigma$ are the initial amplitude and phase of the oscillation, respectively. $\omega_p = \sqrt{(2N_c + N_f)g^2 T^2/18}$ is the plasma frequency, with N_f being the number of flavors for quarks. The dispersion relation Eq.(12) agrees with the leading order result using the hard thermal loops in finite temperature QCD and the classical nature of the hard thermal loops has been investigated extensively by Blaizot and Jancu, etc. [9,10]. As shown by U. Heinz, the eigenwaves satisfying Eq.(12) are always timelike, i.e., $\frac{\omega}{K^{(0)}} > 1$ (we set $\hbar = c = 1$ for convenience). It means that the phase velocity of eigenwaves is bigger than the velocity of light. So these waves can't exchange energy with plasma particles and do not undergo damping in the linear approximation [1,2]. In the long-wavelength region, the dispersion relation Eq.(12) is reduced to the form

$$\omega^2 = \omega_p^2 + \frac{3}{5} K^{(0)2} \quad (14)$$

The following identities will be used to simplify the color current contributed by gluons in the following discussion:

$$\begin{aligned}
f_{abc}f_{abd} &= N_c\delta_{cd}; & if_{abc}G_{bc} &= tr(F_a G); & [F_a, F_b] &= if_{abc}F_c; \\
I_d tr(F_d[F_a, F_b]) &= N_c[I_a, I_b]; & I^d tr(F^d[F^a, [F^b, F^c]]) &= N_c[I^a, [I^b, I^c]]; \\
tr(\{F_a, \{F_b, F_c\}\}F_d) &= 4\delta_{ad}\delta_{cb} + 2\delta_{ab}\delta_{cd} + 2\delta_{ac}\delta_{bd} + N_c d^{ade}d^{bce}
\end{aligned} \tag{15}$$

The second order equations can be discussed in the same manner. Substituting the results in the expression for the second color current by using the distribution functions obtained from the second order kinetic equations, one can find the second order mean field equation is reduced to

$$\begin{aligned}
-\omega^2 \epsilon(\omega, \mathbf{k}^{(0)}) A^{(2)}(k) &= \int \frac{d^3 p}{(2\pi)^3} \left[N_f \frac{d}{dE_p} \left(f^{(0)}(p) + \bar{f}^{(0)}(p) \right) + 2N_c \frac{dG^{(0)}(p)}{dE_p} \right] \\
&\quad \left\{ g^3 \frac{1}{p \cdot k^{(0)} + ip^{00+}} \frac{\mathbf{p} \cdot \mathbf{k}^{(0)}}{K^{(0)}} \sum_{k_1+k_2=k} \frac{\omega_2}{p \cdot k_2 + ip^{00+}} \frac{\mathbf{p} \cdot \mathbf{k}_1^{(0)}}{K_1^{(0)}} \frac{\mathbf{p} \cdot \mathbf{k}_2^{(0)}}{K_2^{(0)}} \right. \\
&\quad \left. [A^{(1)}(k_1), A^{(1)}(k_2)] - g^2 \frac{\mathbf{p} \cdot \mathbf{k}^{(1)}}{p^0} \frac{A^{(1)}(k)}{(p \cdot k^{(0)} + ip^{00+})^2} \frac{(\mathbf{p} \cdot \mathbf{k}^{(0)})^3}{(K^{(0)})^2} \right\} \tag{16}
\end{aligned}$$

Eq.(16) describes the three-wave processes owing to the nonlinear coupling term $A^{(1)}(k_1)A^{(1)}(k_2)$ of two eigenwaves with wave vectors $\mathbf{k}_1$ and $\mathbf{k}_2$ into a secondary wave with wave vector $\mathbf{k}$. However, as pointed out in the Refs. [11, 12], the three-wave processes are forbidden, there will be nonlinear effects of the three-wave processes on wave vectors only in higher order perturbation.

Now let's discuss the third order equations. The third order field equation is

$$\begin{aligned}
-\omega^2 A^{(3)}(k) \frac{k^{(0)h}}{K^{(0)}} + g \sum_{k_1+k_2=k} \frac{k^{(1)i} k_1^{(0)i} k_2^{(0)h}}{K_1^{(0)} K_2^{(0)}} [A^{(1)}(k_1), A^{(1)}(k_2)] \\
+ g^2 \sum_{k_1+k_2=k} \sum_{k_3+k_4=k_2} \frac{k^{(0)i} k_3^{(0)i} k_4^{(0)h}}{K_1^{(0)} K_2^{(0)} K_4^{(0)}} [A^{(1)}(k_1), [A^{(1)}(k_3), A^{(1)}(k_4)]] = j^{(3)h}(k) \tag{17}
\end{aligned}$$

and the third order kinetic equation for quark is

$$p \cdot k^{(0)} f^{(3)}(k, p) - k_i^{(2)} p_i f^{(1)}(k, p) - g p_i \sum_{k_1+k_2=k} [A_i^{(1)}(k_1), f^{(2)}(k_2, p)] - p_i k_i^{(1)} f^{(2)}(k, p)$$

$$\begin{aligned}
& -gp_i \sum_{k_1+k_2=k} [A_i^{(2)}(k_1), f^{(1)}(k_2, p)] + \frac{g}{2} \sum_{k_1+k_2=k} p \cdot k_1^{(0)} \{A_i^{(1)}(k_1), \partial_p^i f^{(2)}(k_2, p)\} \\
& + \frac{g}{2} \sum_{k_1+k_2=k} p \cdot k_1^{(0)} \{A_i^{(2)}(k_1), \partial_p^i f^{(1)}(k_2, p)\} - \frac{g}{2} \sum_{k_1+k_2=k} p_i k_{1i}^{(1)} \{A_j^{(2)}(k_1), \partial_p^j f^{(0)}(k_2, p)\} \\
& + \frac{g}{2} \sum_{k_1+k_2=k} p_i \{A_i^{(1)}(k_1), k_{1\nu}^{(0)} \partial_p^\nu f^{(2)}(k_2, p)\} + \frac{g}{2} \sum_{k_1+k_2=k} p_i \{A_i^{(2)}(k_1), k_{1\nu}^{(0)} \partial_p^\nu f^{(1)}(k_2, p)\} \\
& + \frac{g}{2} \sum_{k_1+k_2=k} p_i \{A_i^{(2)}, k_{1\nu}^{(1)} \partial_p^\nu f^{(0)}(k_2, p)\} + \frac{g}{2} \sum_{k_1+k_2=k} (p \cdot k) \{A_i^{(3)}(k_1), \partial_p^i f^{(0)}(k_2, p)\} \\
& - \frac{g}{2} \sum_{k_1+k_2=k} p_i k_{1i}^{(1)} \{A_j^{(2)}(k_1), \partial_p^j f^{(0)}(k_2, p)\} - \frac{g}{2} \sum_{k_1+k_2=k} p_i k_{1i}^{(2)} \{A_j^{(1)}(k_1), \partial_p^j f^{(0)}(k_2, p)\} \\
& + \frac{g}{2} \sum_{k_1+k_2=k} p_i \{A_i^{(1)}(k_1), k_{1\nu}^{(1)} \partial_p^\nu f^{(1)}(k_2, p)\} + \frac{g}{2} \sum_{k_1+k_2=k} p_i \{A_i^{(1)}(k_1), k_{1\nu}^{(2)} \partial_p^\nu f^{(0)}(k_2, p)\} \\
& + \frac{g}{2} \sum_{k_1+k_2=k} p_i \{A_i^{(3)}(k_1), k_{1\nu}^{(0)} \partial_p^\nu f^{(0)}(k_2, p)\} \\
& - \frac{g^2}{2} \sum_{k_1+k_2=k} \sum_{k_3+k_4=k_1} p_i \{[A_i^{(1)}(k_3), A_j^{(2)}(k_4)], \partial_p^j f^{(0)}(k_2, p)\} \\
& - \frac{g^2}{2} \sum_{k_1+k_2=k} \sum_{k_3+k_4=k_1} p_i \{[A_i^{(1)}(k_3), A_j^{(1)}(k_4)], \partial_p^j f^{(1)}(k_2, p)\} \\
& - \frac{g^2}{2} \sum_{k_1+k_2=k} \sum_{k_3+k_4=k_1} p_i \{[A_i^{(2)}(k_3), A_j^{(1)}(k_4)], \partial_p^j f^{(0)}(k_2, p)\} = 0
\end{aligned} \tag{18}$$

The third order kinetic equations for anti-quarks and gluons are similar except for the changes pointed out in Eq.(8).

Analogously to the electromagnetic plasma theory ^[13], the wave vector $\mathbf{k}$ can be expressed as $\mathbf{k} = \vec{\beta} - i\vec{\alpha}$, where $\beta = |\vec{\beta}|$ is the phase constant and $\alpha = |\vec{\alpha}|$ is damping constant, i.e., the amplitude of the eigenwave will exponentially decrease in the direction of $\vec{\alpha}$. The nonlinear spatial damping rate at this order is defined as

$$\alpha = -Im|\mathbf{k}^{(2)}| = -ImK^{(2)}. \tag{19}$$

As the oscillations are developed from random thermal motions, we have $\langle A^{(1)a}(k) \rangle = 0$, where $\langle \rangle$ means the average with respect to the random phase of the oscillations. So we obtain

$$\begin{aligned}
\langle A^{(1)a}(k) A^{(1)b*}(k') \rangle &= (2\pi)^4 \delta^4(k - k') \delta_a^b \langle A^{(1)2}(k) \rangle_{\mathbf{k}^{(0)}\omega}; \\
\langle A^{(1)2}(k) \rangle_{\mathbf{k}^{(0)}\omega} &= \frac{\pi}{\omega^2} [\delta(\omega - \omega_{\mathbf{k}^{(0)}})] I_{\mathbf{k}^{(0)}}; \quad I_{\mathbf{k}^{(0)}} = \frac{|E_{\mathbf{k}^{(0)}}|}{2V}
\end{aligned} \tag{20}$$

where V is the volume of the plasma; $I_{\mathbf{k}^{(0)}}$ characterizes the total intensity of the fluctuating oscillation with frequency $\omega_{\mathbf{k}}$ and $-\omega_{\mathbf{k}}$. In an equilibrium plasma and the long wavelength limit ($K^{(0)} = 0$) there are $I_{\mathbf{k}^{(0)}} = 4\pi T$. The average of the product of the field potential can be expanded as ^[7,14]

$$\begin{aligned}\langle A^{(1)}(k_1)A^{(1)}(k_2)A^{(1)}(k_3)A^{(1)}(k_4) \rangle &= \langle A^{(1)}(k_1)A^{(1)}(k_2) \rangle \langle A^{(1)}(k_3)A^{(1)}(k_4) \rangle \\ &+ \langle A^{(1)}(k_1)A^{(1)}(k_3) \rangle \langle A^{(1)}(k_2)A^{(1)}(k_4) \rangle \\ &+ \langle A^{(1)}(k_1)A^{(1)}(k_4) \rangle \langle A^{(1)}(k_2)A^{(1)}(k_3) \rangle\end{aligned}\quad (21)$$

By inserting the third order distribution functions Eqs.(18) into Eq.(17), multiplying both sides of the equation by $A^{(1)b}(k')$ and performing the average of the results with respect to the random phase, we have

$$\begin{aligned}-Im \int \frac{d^3p}{E_p(2\pi)^3} \left(\frac{\mathbf{p} \cdot \mathbf{k}^{(0)}}{K^{(0)}} \right)^2 \frac{\mathbf{p} \cdot \mathbf{k}^{(2)}}{(p \cdot k^{(0)} + ip^{00+})^2} \left[N_f \frac{df^{(0)}(p)}{dE_p} + N_c \frac{dG^{(0)}(p)}{dE_p} \right] &= A_1 + A_2; \quad (22) \\ A_1 = Im \left(-\frac{g^2}{\omega_p} \int \frac{d^3p}{(2\pi)^3 E_p} \int \frac{d^4k_1}{(2\pi)^4} \langle A^{(0)2} \rangle_{\mathbf{k}_1^{(0)}\omega} \left(\frac{\omega_1}{K_1^{(0)}} \right)^2 \left(\frac{\omega}{K^{(0)}} \right)^2 \right. \\ &\times \frac{1}{p \cdot (k^{(0)} - k_1^{(0)}) + ip^{00+}} \left(\frac{p^0 \mathbf{k}^{(0)} \cdot \mathbf{k}_1^{(0)}}{p \cdot k^{(0)} + ip^{00+}} - \mathbf{p} \cdot \mathbf{k}^{(0)} \frac{\omega \mathbf{p} \cdot \mathbf{k}_1^{(0)} - p^0 \mathbf{k}^{(0)} \cdot \mathbf{k}_1^{(0)}}{(p \cdot k^{(0)} + ip^{00+})^2} \right) \\ &\times \left\{ \left[\mathbf{k}^{(0)} \cdot \mathbf{k}_1^{(0)} \left(\frac{1}{p \cdot k_1^{(0)} + ip^{00+}} - \frac{1}{p \cdot k^{(0)} + ip^{00+}} \right) + (\mathbf{p} \cdot \mathbf{k}_1^{(0)}) \frac{-\omega_1 \mathbf{p} \cdot \mathbf{k}^{(0)}/p^0 + \mathbf{k}^{(0)} \cdot \mathbf{k}_1^{(0)}}{(\mathbf{p} \cdot \mathbf{k}_1^{(0)} + ip^{00+})^2} \right. \right. \\ &- (\mathbf{p} \cdot \mathbf{k}) \frac{-\omega \mathbf{p} \cdot \mathbf{k}_1^{(0)}/p^0 + \mathbf{k}^{(0)} \cdot \mathbf{k}_1^{(0)}}{(p \cdot k^{(0)} + ip^{00+})^2} \left. \left. \right] \left(\frac{7}{6} N_f \frac{df^{(0)}(p)}{dE_p} + \frac{22}{3} \frac{dG^{(0)}(p)}{dE_p} \right) \right. \\ &+ (\mathbf{p} \cdot \mathbf{k}_1^{(0)}) \frac{(\mathbf{p} \cdot \mathbf{k}^{(0)})}{p^0} \left(\frac{1}{p \cdot k_1^{(0)}} - \frac{1}{p \cdot k^{(0)} + ip^{00+}} \right) \left. \left. \left(\frac{7}{6} N_f \frac{d^2 f^{(0)}(p)}{dE_p^2} + \frac{22}{3} \frac{d^2 G^{(0)}(p)}{dE_p^2} \right) \right] \right\}; \\ A_2 = Im \left\{ \int \frac{d^3p}{(2\pi)^3 E_p} \int \frac{d^4k_1}{(2\pi)^4} \langle A^{(1)2} \rangle_{\mathbf{k}_1^{(0)}\omega} \left(N_f \frac{df^{(0)}}{dE_p} + N_c \frac{dG^{(0)}(p)}{dE_p} \right) \right. \\ &\times \frac{1}{p \cdot k^{(0)} + ip^{00+}} \times \left[\frac{-6g^2}{\omega_p} \left(\frac{\mathbf{p} \cdot \mathbf{k}^{(0)}}{K^{(0)}} \right)^2 \left(\frac{\mathbf{p} \cdot \mathbf{k}^{(0)}}{K_1^{(0)}} \right)^2 \frac{1}{p \cdot (k^{(0)} - k_1^{(0)}) + ip^{00+}} \right. \\ &\times \left(\frac{\omega_1}{p \cdot k_1 + ip^{00+}} - \frac{\omega}{p \cdot k^{(0)} + ip^{00+}} \right) + \frac{12g^4}{\omega_p} \left(\frac{\mathbf{p} \cdot \mathbf{k}^{(0)}}{K^{(0)}} \right) \left(\frac{\mathbf{p} \cdot \mathbf{k}_1^{(0)}}{K_1^{(0)}} \right) \left(\frac{\mathbf{p} \cdot (\mathbf{k}^{(0)} - \mathbf{k}_1^{(0)})}{|\mathbf{k}^{(0)} - \mathbf{k}_1^{(0)}|} \right) \\ &\times \left. \left(\frac{\omega_1}{p \cdot k_1 + ip^{00+}} - \frac{\omega}{p \cdot k^{(0)} + ip^{00+}} \right) \times \int \frac{d^3p'}{(2\pi)^3 E'_p} \left(N_f \frac{df^{(0)}(p')}{dE'_p} + N_c \frac{dG^{(0)}(p')}{dE'_p} \right) \right\}\end{aligned}$$

$$\begin{aligned} & \times \left(\frac{\mathbf{p}' \cdot (\mathbf{k}^{(0)} - \mathbf{k}_1^{(0)})}{|\mathbf{k}^{(0)} - \mathbf{k}_1^{(0)}|} \right) \left(\frac{\mathbf{p}' \cdot \mathbf{k}^{(0)}}{K^{(0)}} \right) \left(\frac{\mathbf{p}' \cdot \mathbf{k}_1^{(0)}}{K_1^{(0)}} \right) \frac{1}{(\omega - \omega_1)^2} \frac{1}{\epsilon(\omega - \omega_1, \mathbf{k}^{(0)} - \mathbf{k}_1^{(0)})} \\ & \times \frac{1}{p' \cdot (k^{(0)} - k_1^{(0)}) + ip^{00+}} \left(\frac{\omega_1}{p' \cdot k_1^{(0)} + ip^{00+}} - \frac{\omega}{p' \cdot k^{(0)} + ip^{00+}} \right) \Bigg] \Bigg\} \end{aligned}$$

To get the numerical result of the nonlinear spatial damping rate, we perform the integrals in cylindrical coordinates and in the local rest frame of the plasma particle.

At first the left hand side of Eq.(22) can be simplified by averaging over all directions of $\mathbf{k}^{(2)}$ and selecting the direction of $\mathbf{k}^{(0)}$ as the direction of the polar axis, i.e., we have

$$\begin{aligned} & -Im \int \frac{d^3p}{E_p(2\pi)^3} \frac{(\mathbf{p} \cdot \mathbf{k}^{(0)})^2}{(K^{(0)})^2} \frac{\mathbf{p} \cdot \mathbf{k}^{(2)}}{(p \cdot k^{(0)} + ip^{00+})^2} \left(N_f \frac{d}{dE_p} f^{(0)}(p) + N_c \frac{dG^{(0)}(p)}{dE_p} \right) \\ & = -Im \langle K^2 \rangle \frac{1}{3\omega_p^2} \int \frac{d^3p}{E_p(2\pi)^3} \cos^2 \phi \left(N_f \frac{d}{dE_p} f^{(0)}(p) + N_c \frac{dG^{(0)}(p)}{dE_p} \right) \end{aligned} \quad (23)$$

where ϕ is the angle between $\mathbf{p}$ and $\mathbf{k}^{(0)}$. $\langle \rangle$ means the average over all directions of $\mathbf{k}^{(2)}$ [14,15].

Now we can see that $Im \langle K^{(2)} \rangle$ can be extracted from the left hand side of Eq.(22), and the numerical value of $Im \langle K^{(2)} \rangle$ is determined by A_1 and A_2 . Before performing the integrals in A_1 and A_2 , we analyze the mechanism of the nonlinear spatial damping. The important relation $p^0/(p \cdot k' + ip^{00+}) = P [1/(\omega' - \mathbf{v} \cdot \mathbf{k}') - i\pi\delta(\omega' - \mathbf{v} \cdot \mathbf{k}')] is very useful in our discussion, where $\mathbf{v} = \mathbf{p}/p^0$ is the velocity of particle and P stand for the principal value of the function. We can see that $Im \langle K^{(2)} \rangle$ is not vanishing only when at least one of the imaginary parts of $1/(p \cdot k^{(0)} + ip^{00+})$, $1/(p \cdot k_1^{(0)} + ip^{00+})$ and $1/(p \cdot (k^{(0)} - k_1^{(0)}) + ip^{00+})$ is not zero. As is shown in the linear approximation, the color eigenwaves are timelike, i.e., the phase velocity of waves can't approach the velocity of particle and the imaginary parts of $1/(p \cdot k^{(0)} + ip^{00+})$ and $1/(p \cdot k_1^{(0)} + ip^{00+})$ are zero. The term $1/(p \cdot (k^{(0)} - k_1^{(0)}) + ip^{00+})$ appearing in A_1 and A_2 are very crucial, it describes that the eigenwaves with wave vectors $\mathbf{k}^{(0)}$ and $\mathbf{k}_1^{(0)}$ may produce the secondary waves with wave vectors $\mathbf{k}^{(0)} - \mathbf{k}_1^{(0)}$ through the nonlinear interactions. Even if the eigenwaves are timelike, these secondary waves may be spacelike, i.e., their phase velocity $\frac{\omega - \omega_1}{\mathbf{k}^{(0)} - \mathbf{k}_1^{(0)}}$ may be smaller than the velocity of light and can approach the velocity of particle. So the imaginary part of $\delta((\omega - \omega_1) - \mathbf{v} \cdot (\mathbf{k}^{(0)} - \mathbf{k}_1^{(0)}))$$

may not be zero and the secondary waves may exchange energy with plasma particles and be damped by the particles. It should be emphasized that though the mechanism is analogous to the mechanism of nonlinear Landau damping in electromagnetic plasmas, the nonlinear coupling of the waves takes place through the nonlinear relations $[A^{(1)}(k_1), A^{(1)}(k_2)]$ etc. in Eq.(17), i.e., it describes the nonlinear and non-Abelian characteristics of QGP.

The similar integral to A_1 and A_2 has been discussed in Ref. [4], by taking only the leading order in g and taking the long-wavelength limit, we find

$$A_1 + A_2 \sim 0.42T. \quad (24)$$

After finishing the integral of the right hand side of Eq.(23), and from Eq.(24), we obtain the nonlinear spatial damping rate for a pure gluon gas in the long wavelength limit

$$\alpha = -Im\langle K^{(2)} \rangle \sim 2.52g^2T \quad (25)$$

Now we summarize briefly. In this paper, we use the derivative expansion method to study the nonlinear and the non-Abelian effects given by the kinetic equations of QGP. By solving the equations to the third order, the numerical result of the spatial damping rate α for the secondary waves in the long wavelength limit is obtained. This result indicates that the amplitude of the secondary waves in QGP decrease exponentially, and its damping rate is $2.52g^2T$ Neper approximately. It makes us have a concrete understanding about the spatial traveling of color waves in QGP.

REFERENCES

- [1] H.-Th. Elze, Z. Phys. C**38**,211(1988).
- [2] U. Heinz, Ann. Phys. (N.Y.) **168**,148(1986)
- [3] Mrówczyński, Phys. Rev. D**39**,1940(1989).
- [4] Zhang Xiaofei and Li Jiarong, Phys. Rev. C**52**,964(1995).
- [5] T. H. Stix, *Waves in Plasmas*(American Institute of Physics, New York, 1992)
- [6] H.-Th. Elze and U. Heinz,Phys. Rep. **183**,81(1989).
- [7] A. G. Sitenko, *Fluctuations and Nonlinear Wave Interactions in Plasmas*(Pergamon, Oxford, 1990).
- [8] A. Jeffrey and T. Kawahara, *Asymptotic Method in Nonlinear Wave theory*(Pitman, London ,1982).
- [9] J.-P. Blaizot and E. Jancu, Phys.Rev.Lett.**70**,3376(1993); Nucl.Phys.B**421**,565(1994).
- [10] R. Jackiw, Q. Liu and C. Lucchesi, Phys. Rev.D**49**,6787(1994);
P. F. Kelly, Q. Liu, C. Lucchesi and C. Manuel, Phys. Rev. Lett. **72**,3461(1994).
- [11] B. B. Kadomtsev, *Plasma Turbulence*(Academic, New York, 1965)
- [12] J. Weiland and H. Wilhelmsson, *Coherent Nonlinear Interaction of Waves in Plasma*(Pergamon, Oxford, 1976).
- [13] H. G. Booker,*Electromagnetic and Hydromagnetic Waves in a Cold Magnetic Plasma*
(Phil. Trans. R. Soc. Lond., A**280**,1975)
- [14] L. D. Landau and E. M. Lifshitz, *Electrodynamics of Continuous Media*(Pergamon, New York, 1960).
- [15] E. M. Lifshitz and L. P. Pitaevskii,*Physical Kinetics*(Pergamon, Oxford,1981).